

 View PDFAccess through **your institution**[Purchase PDF](#)

South African Journal of Botany

Volume 135, December 2020, Pages 80-86

# Effect of the medium composition on the asymbiotic germination and *in vitro* development of the *Laeliocattleya* hybrid

Seir Antonio Salazar Mercado <sup>a</sup>  , Edison Alexander Botello Delgado <sup>b</sup>  [Show more](#)  Outline |  Share  Cite<https://doi.org/10.1016/j.sajb.2020.08.011>[Get rights and content](#)

## Highlights

- Preconditioning treatment influence *xLaeliocattleya* hybrid seeds viability, tested by tetrazolium staining.
- The immersion in 10% sucrose and distilled water for 10 min improved the detection of viable seeds.
- The best germination percentage was achieved in the culture medium supplemented with pineapple juice ( $56 \pm 2.8$ ).
- The MS medium enriched with pineapple juice proved to be efficient in the seeds germination and development of *xLaeliocattleya*.
- The use of organic supplements in the culture medium could replace synthetic hormones.

FEEDBACK 

 View PDFAccess through **your institution**[Purchase PDF](#)

enhance the tetrazolium test. Initially, the indehiscent capsules were collected in a nursery in Bochalema-Norte de Santander. From them, seeds were obtained for the preconditioning and germination tests. One portion was subjected to different precondition treatments: distilled water, sodium hypochlorite (0.5–1.0%) and sucrose (10%). Afterwards, seeds were rinsed and exposed to different concentrations and exposure periods of tetrazolium. Regarding *in vitro* cultivation, the commercial medium Orchid Seed Sowing Medium, Murashige & Skoog (MS), and the MS supplemented with coconut water (MSC), pineapple juice (MSP) and gibberellic acid (MSG3) were evaluated. The highest percentage of viability was obtained by subjecting the seeds to preconditioning with 10% sucrose and keeping them for 24 h in a 0.5% tetrazolium solution. With respect to the MS media used, those enriched with organic supplements proved to be superior to the other treatments since the highest results were achieved both in asymbiotic germination ( $56 \pm 2.8\%$ ) and in seedlings formation ( $25.8 \pm 0.8$ ) with the MSP medium, followed by the MSC medium. In this way, a viable and low-cost alternative is offered for the commercial propagation of *xLaeliocattleya* Richard Muller.

[< Previous](#)[Next >](#)

## Keywords

Development; *In vitro* germination; Hybrid; Viability

[Recommended articles](#)[Citing articles \(0\)](#)[View full text](#)

© 2020 SAAB. Published by Elsevier B.V. All rights reserved.



Copyright © 2021 Elsevier B.V. or its licensors or contributors.  
ScienceDirect® is a registered trademark of Elsevier B.V.

[FEEDBACK](#) 



View PDF

Access through **your institution**

[Purchase PDF](#)

---

FEEDBACK 