



ScienceDirect

 View PDFAccess through **your institution**[Purchase PDF](#)

Science of The Total Environment

Volume 700, 15 January 2020, 134452

Cytotoxic evaluation of glyphosate, using *Allium cepa* L. as bioindicator

Seir Antonio Salazar Mercado ^a  , Jesús David Quintero Caleño ^b 

^a Department of Biology, Universidad Francisco de Paula Santander, Avenida Gran Colombia No. 12E-96B Colsag, San José de Cúcuta, Colombia

^b Department of Agricultural Sciences, Universidad Francisco de Paula Santander, Colombia

Received 1 August 2019, Revised 7 September 2019, Accepted 13 September 2019, Available online 4 October 2019.

Editor: Lotfi Aleya

[Show less](#)  Outline |  Share  Cite<https://doi.org/10.1016/j.scitotenv.2019.134452>[Get rights and content](#)

Highlights

- The use of glyphosate caused permanent DNA damage.


FEEDBACK 

- The 30 mg L⁻¹ treatment caused an average frequency of 170 elongated nuclei.
- The exposition to 15 and 30 mg L⁻¹ glyphosate caused an average frequency of 58 micronuclei.

Abstract

Glyphosate is a chemical compound used mainly as a broad spectrum herbicide, it is recognized for its proven effectiveness and easy handling. It represents more than 60% of the world market of non-selective herbicides and is used in both agricultural fields and family gardens. The present study was designed to test the cytogenotoxic potential of glyphosate using the *Allium cepa* test as toxicity bioindicator. Consequently, bulbs of *A. cepa* were exposed to different concentrations of glyphosate (5, 10, 15, 25 and 30 mgL⁻¹) and a control (deionized water), for 72 h; root development was also studied in this period of time. The cytogenotoxic potential of glyphosate was determined by calculating the mitotic index (MI), cellular anomalies (CA) and registering the roots longitudinal growth at 24, 48 and 72 h. Regarding root development, a greater growth was observed in the control treatment in the three measurement times. The mitotic phases analysis, determined that the higher the concentration, the lower the mitotic index, in addition the inhibition of the telophase Mitotic Index (TMI) was observed in any of the concentrations. The results indicate that the exposure to glyphosate of *A. cepa* meristematic cells induces diverse types of chromosomal anomalies, such as micronuclei (MN), chromosome breaking (CB), nuclear notch (Nn), among others. Therefore, in demonstrates that glyphosate has a highly cytogenotoxic effect for any of the concentrations used.

Graphical abstract



[Download : Download high-res image \(315KB\)](#)

[Download : Download full-size image](#)

[Previous](#)

[Next](#)

FEEDBACK

Keywords

Cellular anomalies; Mitosis; Chromosome; *Allium cepa*; Cytogenotoxic; DNA

[Recommended articles](#)

[Citing articles \(14\)](#)

[View full text](#)

© 2019 Elsevier B.V. All rights reserved.



[About ScienceDirect](#)

[Remote access](#)

[Shopping cart](#)

[Advertise](#)

[Contact and support](#)

[Terms and conditions](#)

[Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the **use of cookies**.

Copyright © 2021 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

ScienceDirect® is a registered trademark of Elsevier B.V.

